

OFFICE OF NAVAL RESEARCH  
CONTRACT N00014-88-C-0118

TECHNICAL REPORT 92-03

RED BLOOD CELL VOLUME, PLASMA VOLUME AND TOTAL BLOOD VOLUME  
IN HEALTHY ELDERLY MEN AND WOMEN AGED 64 TO 100

BY

C.R. VALERI, L.E. PIVACEK, H. SIEBENS, and M.D. ALTSCHULE

NAVAL BLOOD RESEARCH LABORATORY  
BOSTON UNIVERSITY SCHOOL OF MEDICINE  
615 ALBANY STREET  
BOSTON, MA 02118

6 MAY 1992

Reproduction in whole or in part is permitted for any  
purpose of the United States Government.

Distribution of this report is unlimited.

DTIC QUALITY INSPECTED 1

19990225060

## UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NBRL, BUSM 92-03	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) RED BLOOD CELL VOLUME, PLASMA VOLUME AND TOTAL BLOOD VOLUME IN HEALTHY ELDERLY MEN AND WOMEN AGED 64 TO 100		5. TYPE OF REPORT & PERIOD COVERED Technical Report
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) C. Robert Valeri, Linda E. Pivacek, Hiliary Siebens, and Mark D. Altschule		8. CONTRACT OR GRANT NUMBER(s) N00014-88-C-0118
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Blood Research Laboratory Boston University School of Medicine 615 Albany St., Boston, MA 02118		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS Naval Medical Research and Development Command Bethesda, MD 20814		12. REPORT DATE 6 May 1992
		13. NUMBER OF PAGES 29
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Bureau of Medicine and Surgery Department of the Navy Washington, D.C. 20372		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release and sale. Distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Red blood cell volume                      Blood Plasma volume                                Hematocrit Total blood volume Red blood cells		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) In healthy men and women aged 64 to 100 years of age, red blood cell volume was measured using <sup>51</sup> Cr-labeled fresh autologous red blood cells, and plasma volume was measured using <sup>125</sup> I albumin. These measurements were correlated to body surface area (BSA) and body weight (BW). In these elderly healthy male and female subjects, red cell volumes were 11 to 23% lower than, plasma volumes were similar to, and total blood volumes were slightly lower than values previously reported in younger subjects. The		

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

red blood cell volume showed no correlation with the peripheral venous hematocrit or the red blood cell 2,3 DPG, ATP, or P50.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

ABSTRACT: In healthy men and women aged 64 to 100 years of age, red blood cell volume was measured using  $^{51}\text{Cr}$ -labeled fresh autologous red blood cells, and plasma volume was measured using  $^{125}\text{I}$  albumin. These measurements were correlated to body surface area (BSA) and body weight (BW).

In these elderly healthy male and female subjects, red cell volumes were 11 to 23% lower than, plasma volumes were similar to, and total blood volumes were slightly lower than values previously reported in younger subjects. The red blood cell volume showed no correlation with the peripheral venous hematocrit or the red blood cell 2,3 DPG, ATP, or p50.

We report the red blood cell, plasma and total blood volumes in relation to body surface area (BSA) and body weight (BW), and report regression equations used to estimate red blood cell, plasma and total blood volumes.

INTRODUCTION: Our study of healthy men and women 64 to 100 years of age was done to determine whether individuals in this age group have blood volume measurements that are different from those of younger individuals<sup>1-10</sup>. We used  $^{51}\text{Cr}$ -labeled autologous red cells to measure red blood cell volume, and  $^{125}\text{I}$  albumin to measure plasma volume. We also measured red blood cell ATP, DPG, and p50 levels.

METHODS: The 51 healthy elderly volunteers (32 females and 19 males) who participated in the study signed informed consent forms in accordance with the Institutional Review Board for Human Research at Boston University Medical

Center. Participants were interviewed and were medically examined to determine their health status and use of medications. Each subject was screened for cardiovascular, renal, liver, and metabolic diseases. All participants were physically active and in good health.

The subjects were maintained in a supine position for 30 minutes prior to infusion of the  $^{51}\text{Cr}$ -labeled red blood cells and the  $^{125}\text{I}$  albumin. To measure red blood cell volume, a 20 milliliter (ml) blood sample was collected, incubated at 37 degrees Centigrade for 30 minutes with 11.5 microcuries of  $^{51}\text{Cr}$  disodium chromate (ER Squibb & Sons). A 10 ml volume of  $^{51}\text{Cr}$ -labeled fresh autologous red blood cells containing 5 microcuries was reinfused. To measure plasma volume, a solution of 1.5 microcuries of  $^{125}\text{I}$  human albumin in isotonic saline was prepared, and a 5 ml volume of the solution containing 0.5 microcuries was infused.

Venous blood samples were obtained before infusion and 20 and 30 minutes after infusion for measurements of radioactivity in the blood and plasma. Red blood cell volume (RCV) was calculated from the total  $^{51}\text{Cr}$  radioactivity associated with the injected red blood cells and  $^{51}\text{Cr}$  radioactivity associated with the red blood cells 20 and 30 minutes after infusion. Plasma volume (PV) was calculated from the total injected  $^{125}\text{I}$  radioactivity and the mean plasma  $^{125}\text{I}$  radioactivity 20 and 30 minutes following infusion. The total blood volume was calculated

as the sum of the  $^{51}\text{Cr}$  red blood cell volume and the  $^{125}\text{I}$  albumin plasma volume.

The peripheral venous microhematocrit was measured before and after the infusions of the  $^{51}\text{Cr}$  labeled red blood cells and the  $^{125}\text{I}$  albumin. The total body hematocrit was calculated from the red cell volume divided by the total blood volume,  $\text{RCV} / (\text{RCV} + \text{PV})$ , and the f-value was calculated from the total body hematocrit and the peripheral venous hematocrit <sup>11,12,13</sup>.

At the time that the red blood cell and plasma volumes were being measured, peripheral venous blood was collected to measure red blood cell 2,3 DPG, ATP, and p50 levels. Red blood cell adenosine triphosphate (ATP) and 2,3 diphosphoglycerate (2,3 DPG) levels were measured fluorometrically<sup>14,15</sup>. The red blood cell p50 was measured in washed red blood cells in a phosphate buffer solution, pH 7.2 at a temperature of +37C and a pCO<sub>2</sub> tension of 0 mm Hg (Hemoscan Oxygen Dissociation Analyzer, American Instruments Co., Silver Spring, MD)<sup>16</sup>. The pO<sub>2</sub> tension at which 50 percent of the hemoglobin was saturated with oxygen is reported as the p50 value.

Statistical analyses were performed using Statistical Analysis Systems (SAS) licensed to Boston University, and included means, standard deviations, and linear regression and correlation.

RESULTS: Table 1 reports the age, height, body weight (BW), body surface area (BSA), and peripheral venous hematocrit of

the 19 men and 32 women in the study. The mean age of the men was 73, with a range of 64 to 100; the mean age of the women was 74, with a range of 65 to 93. No significant correlation was seen between the age of the male and female subjects and their red blood cell, plasma, or total blood volume.

The mean  $^{51}\text{Cr}$  red cell volume (RCV) for the men was 1662 ml, and when this measurement was related to body indices, the value was 889 ml/m<sup>2</sup> for body surface area (BSA) and 22.6 ml/kg for body weight (BW) (Table 2). The regression equation relating the red cell volume (ml) to body surface area (m<sup>2</sup>) was  $\text{RCV} = 1761(\text{m}^2) - 1608$ , and to weight (kg) was  $\text{RCV} = 25.1 (\text{kg}) - 183$  (Table 3). The body surface area (BSA) accounted for 75% of the variance in red cell volume with a correlation coefficient of  $r=0.868$ , and the body weight (BW) accounted for 72% of the variance with a correlation coefficient of  $r=0.846$  (Figures 1 and 2).

In the male subjects, mean plasma volume measured by  $^{125}\text{I}$  albumin was 3029 ml, 1634 ml/m<sup>2</sup> BSA, and 41.9 ml/kg BW (Table 2). The regression equation relating the plasma volume to body surface area (BSA) was  $\text{PV} = 1995(\text{m}^2) - 667$ , and to weight was  $\text{PV} = 24.3(\text{kg}) + 1255$  (Table 3). The BSA accounted for 43% of the variance in plasma volume with a correlation coefficient of  $r=0.657$ , and the BW accounted for 30% of the variance with a correlation coefficient of  $r=0.549$  (Figures 1 and 2).

The male subjects exhibited a mean total blood volume (TBV) of 4693 ml, 2626 ml/m<sup>2</sup> BSA and 64.6 ml/kg BW (Table 2). The regression equation relating the total blood volume to body surface area was  $TBV = 3809(m^2) - 2362$ , and to weight was  $TBV = 49.9(kg) + 1044$  (Table 3). The BSA accounted for 67% of the variance in blood volume with a correlation coefficient of  $r=0.818$ , and the BW accounted for 54% of the variance with a correlation coefficient of  $r=0.736$  (Figures 1 and 2).

In the female subjects, the mean red cell volume was 1185 ml, and the red blood cell volume was 725 ml/m<sup>2</sup> BSA and 19.3 ml/kg BW (Table 2). The regression equation relating the red blood cell volume to body surface area was  $RCV = 716(m^2) + 14$ , and to weight was  $RCV = 9.7(kg) + 573$ , (Table 3). The BSA accounted for 46% of the variance in red cell volume with a correlation coefficient of  $r=0.681$ , and the BW accounted for 49% of the variance with a correlation coefficient of  $r=0.699$  (Figures 3 and 4).

The mean <sup>125</sup>I albumin plasma volume in the female subjects was 2310 ml, 1424 ml/m<sup>2</sup> BSA and 38.3 ml/kg BW (Table 2). The regression equation relating the plasma volume to body surface area was  $PV = 925(m^2) + 802$ , and to weight was  $PV = 10.9(kg) + 1630$  (Table 3). The BSA accounted for 30% of the variance in plasma volume with a correlation coefficient of  $r=0.553$ , and the BW accounted for 23% of the variance with a correlation coefficient of  $r=0.484$  (Figures 3 and 4).



The female subjects exhibited a mean total blood volume of 3482 ml, 2144 ml/m<sup>2</sup>, and 57.6 ml/kg (Table 2). The regression equation relating the blood volume to body surface area was  $TBV = 1591(m^2) + 889$ , and to weight was  $TBV = 19.9(kg) + 2247$  (Table 3). The BSA accounted for 48% of the variance in blood volume with a correlation coefficient of  $r=0.696$ , and the BW accounted for 41% of the variance with a correlation coefficient of  $r=0.642$  (Figures 3 and 4).

The mean f-value in the men was 0.88 with a range of 0.81 to 0.96, and in the women was 0.88 with a range of 0.77 to 1.05 (Table 2).

Table 4 reports the red blood cell volume, plasma volume and total blood volume data from previous studies and from our present study of elderly men and women. The red cell volumes related to body size indices were 11 to 23% lower in the elderly men and women in our study than in the younger subjects in previous studies<sup>3,4,5,7,10</sup>. The plasma volumes in our elderly subjects were within  $\pm 5\%$  those observed in the younger individuals in the previous studies, and blood volumes were 3 to 8% lower<sup>5,7,10</sup>. Data from other studies in subjects over the age of 60 showed plasma volumes within  $\pm 5\%$  of our values in one study<sup>8</sup> and in another study 12% lower than our values for men and 16% lower for women<sup>9</sup>. A third study reports red blood cell volumes 13% higher than our values<sup>10</sup>.

In our study reported here, the age of the subjects did not correlate significantly with the red blood cell 2,3

DPG, ATP or p50 level (Figure 5). Nor was there any significant correlation between peripheral venous hematocrit in the elderly men and women and the red blood cell ATP, DPG, or p50 level (Figures 6, 7 and 8). The red blood cell volume in these subjects did not significantly correlate with red blood cell ATP, 2,3 DPG, or p50 level (Figures 9, 10, 11, 12, 13, and 14).

DISCUSSION: The purpose of our study was to measure red blood cell, plasma, and total blood volumes in healthy elderly men and women aged 64 to 100. We wanted to determine a nomogram from which to predict the red blood cell, plasma and total blood volumes from body surface area and body weight. Red blood cell, plasma and total blood volumes were found to correlate with both body surface area (BSA) and body weight (BW), a finding similar to that reported by other investigators who studied younger subjects 1-10. In our study, the regression equations were slightly more accurate for body surface area (BSA) than for body weight (BW).

The elderly subjects in our study exhibited red blood cell volumes that were consistently lower than those measured in younger subjects using  $^{51}\text{Cr}$  labeled red blood cells 3,4,5,7,10. Plasma volumes were similar to those seen in younger men and women in other studies, and total blood volumes were slightly lower 5,7. The reductions in red blood cell volume in our elderly subjects were not

associated with either increases in plasma volume or decreases in peripheral venous hematocrit <sup>12,13</sup>.

We did not measure lean body mass in our elderly subjects as a means of assessing the correlation between red blood cell volume and the lean body mass. However a study of younger subjects has shown that lean body mass correlated with red blood cell volume<sup>3</sup> and with red blood cell, plasma and total blood volumes<sup>7</sup>.


Plasma volumes that were measured using <sup>125</sup>I albumin and related to the body surface area (BSA) were similar to values seen when T-1824 dye was used<sup>8</sup>. The plasma volumes related to body weight (BW) that were 12% to 16% higher in our study than those reported by Sklaroff<sup>9</sup> may have been due to the fact that our subjects were living at home and were physically active, whereas his subjects were residents in a home for the aged. The subjects in the study by Schmidt, et al<sup>8</sup> were custodial patients in a state hospital.

The normal 2,3 DPG levels and significant reductions in red blood cell volume in our subjects suggest an adaptive red blood cell volume deficiency<sup>17</sup>. The absence of an increase in red blood cell 2,3 DPG accompanied by a significant deficit in red blood cell volume suggests that these elderly patients had an adequate oxygen supply to the tissues. The deficiency in red blood cell volume in our elderly subjects was consistent with an adaptive and not a pathologic deficiency in red blood cell volume.

**FIGURE 1**

RELATIONS BETWEEN BODY SURFACE AREA (BSA) AND RED BLOOD  
CELL, PLASMA AND BLOOD VOLUMES IN ELDERLY MEN


# RELATIONS BETWEEN BODY SURFACE AREA AND RED BLOOD CELL, PLASMA AND BLOOD VOLUMES



**FIGURE 2**

RELATIONS BETWEEN BODY WEIGHT (BW) AND RED BLOOD CELL,  
PLASMA AND BLOOD VOLUMES IN ELDERLY MEN

# RELATIONS BETWEEN BODY WEIGHT AND RED BLOOD CELL, PLASMA AND BLOOD VOLUMES




**FIGURE 3**

RELATIONS BETWEEN BODY SURFACE AREA (BSA) AND RED BLOOD  
CELL, PLASMA AND BLOOD VOLUMES IN ELDERLY WOMEN



# RELATIONS BETWEEN BODY SURFACE AREA AND RED BLOOD CELL, PLASMA AND BLOOD VOLUMES




Body surface area (m²)

- Red blood cell volume
- △ Plasma volume
- ▲ Total blood volume

**FIGURE 4**

RELATIONS BETWEEN WEIGHT (WT) AND RED BLOOD CELL, PLASMA AND  
BLOOD VOLUMES IN ELDERLY WOMEN


# RELATIONS BETWEEN BODY WEIGHT AND RED BLOOD CELL, PLASMA AND BLOOD VOLUMES



**FIGURE 5**

RELATIONS BETWEEN AGE AND RED BLOOD CELL 2,3 DPG, ATP, AND  
P50 LEVELS IN ELDERLY MEN AND WOMEN


# RELATIONS BETWEEN AGE & RED BLOOD CELL 2,3DPG, ATP & P50 IN ELDERLY MEN & WOMEN



**FIGURE 6**

RELATION BETWEEN PERIPHERAL VENOUS HEMATOCRIT AND RED BLOOD  
CELL ATP LEVEL IN ELDERLY MEN AND WOMEN

RELATION BETWEEN PERIPHERAL VENOUS HCT &  
RED BLOOD CELL ATP IN ELDERLY MEN AND WOMEN




**FIGURE 7**

RELATION BETWEEN PERIPHERAL VENOUS HEMATOCRIT AND RED BLOOD  
CELL 2,3 DPG IN ELDERLY MEN AND WOMEN




RELATION BETWEEN PERIPHERAL VENOUS HCT &  
RED BLOOD CELL 2,3DPG IN ELDERLY MEN AND WOMEN



**FIGURE 8**

RELATION BETWEEN PERIPHERAL VENOUS HEMATOCRIT AND RED BLOOD  
CELL P50 IN ELDERLY MEN AND WOMEN


RELATION BETWEEN PERIPHERAL VENOUS HCT &  
RED BLOOD CELL P50 IN ELDERLY MEN & WOMEN



**FIGURE 9**

RELATION BETWEEN RED BLOOD CELL VOLUME AND RED BLOOD CELL  
ATP LEVELS IN ELDERLY MEN


RELATION BETWEEN RED BLOOD CELL VOLUME &  
RED BLOOD CELL ATP IN ELDERLY MEN



**FIGURE 10**

RELATION BETWEEN RED BLOOD CELL VOLUME AND RED BLOOD CELL  
2,3 DPG LEVEL IN ELDERLY MEN

RELATION BETWEEN RED BLOOD CELL VOLUME &  
RED BLOOD CELL 2,3DPG IN ELDERLY MEN




**FIGURE 11**

RELATION BETWEEN RED BLOOD CELL VOLUME AND RED BLOOD CELL  
P50 LEVELS IN ELDERLY MEN




RELATION BETWEEN RED BLOOD CELL VOLUME &  
RED BLOOD CELL P50 IN ELDERLY MEN



**FIGURE 12**

RELATION BETWEEN RED BLOOD CELL VOLUME AND RED BLOOD CELL  
ATP LEVELS IN ELDERLY WOMEN


RELATION BETWEEN RED BLOOD CELL VOLUME  
& RED BLOOD CELL ATP IN ELDERLY WOMEN



**FIGURE 13**

RELATION BETWEEN RED BLOOD CELL VOLUME AND RED BLOOD CELL  
2,3 DPG LEVELS IN ELDERLY WOMEN

RELATION BETWEEN RED BLOOD CELL VOLUME &  
RED BLOOD CELL 2,3DPG IN ELDERLY WOMEN



**FIGURE 14**

RELATION BETWEEN RED BLOOD CELL VOLUME AND RED BLOOD CELL  
P50 LEVELS IN ELDERLY WOMEN

RELATION BETWEEN RED BLOOD CELL VOLUME &  
RED BLOOD CELL P50 IN ELDERLY WOMEN




Table 1

## HEALTHY ELDERLY VOLUNTEERS

	Age	Height (inches)	Weight (kg)	BSA (m <sup>2</sup> )	Peripheral Venous Hematocrit (%)
<u>Males</u>					
Mean:	73.4	67.5	73.4	1.86	40.0
SD:	9.2	3.4	10.9	0.16	3.4
n:	19	19	19	19	19
Minimum:	64.0	62.0	57.7	1.65	34.0
Maximum:	100.0	74.5	94.5	2.13	46.5
<u>Females</u>					
Mean:	73.8	62.4	62.7	1.63	38.2
SD:	6.9	2.7	13.0	0.17	2.2
n:	32	32	32	32	32
Minimum:	65.0	56.0	37.3	1.22	35.0
Maximum:	93.0	68.0	91.8	2.01	42.4



Table 2

RED BLOOD CELL VOLUMES MEASURED USING <sup>51</sup>CR LABELED  
AUTOLOGOUS RED BLOOD CELLS AND PLASMA VOLUMES MEASURED USING  
<sup>125</sup>I ALBUMIN IN HEALTHY ELDERLY VOLUNTEERS

		<u>Males</u>	<u>Females</u>
51Cr Red Blood Cell Volume (ml)	Mean:	1662	1185
	SD:	325	181
	n:	19	32
125I Albumin Plasma Volume (ml)	Mean:	3029	2310
	SD:	480	289
	n:	18	31
Total Blood Volume (ml)	Mean:	4693	3482
	SD:	757	395
	n:	18	31
51Cr Red Blood Cell Volume (ml/M2 BSA)	Mean:	889	725
	SD:	114	79
	n:	10	32
125I Plasma Volume (ml/M2 BSA)	Mean:	1634	1424
	SD:	201	161
	n:	18	31
Total Blood Volume (ml/M2 BSA)	Mean:	2626	2144
	SD:	251	186
	n:	18	31
51Cr Red Blood Cell Volume (ml/kg)	Mean:	22.6	19.3
	SD:	2.4	2.9
	n:	19	32
125 I Albumin Plasma Volume (ml/kg)	Mean:	41.9	38.3
	SD:	6.5	7.5
	n:	18	31
Total Blood Volume (ml/kg)	Mean:	64.6	57.6
	SD:	7.4	9.8
	n:	18	31
Total Body Hematocrit (%)	Mean:	35.4	33.5
	SD:	3.7	3.2
	n:	18	32
F Value	Mean:	0.88	0.88
	SD:	0.05	0.06
	n:	18	32

TABLE 3

REGRESSION EQUATIONS RELATING RED BLOOD CELL, PLASMA AND  
TOTAL BLOOD VOLUMES TO BODY SURFACE AREA AND BODY WEIGHT IN  
THE ELDERLY MEN AND WOMEN

	<u>Men</u>	<u>Women</u>
Red Blood Cell Volume, ml	$= 1761 (m^2) - 1608$ $25.1 (kg) - 183$	$716 (m^2) + 14$ $9.7 (kg) + 573$
Plasma Volume, ml	$= 1995 (m^2) - 667$ $24.3 (kg) + 1255$	$925 (m^2) + 802$ $10.9 (kg) + 1630$
Total Blood Volume, ml	$= 3809 (m^2) - 2362$ $49.9 (kg) + 1044$	$1591 (m^2) + 889$ $19.9 (kg) + 2247$

Study	Age of Subject	Method	MEN						WOMEN					
			RCV/PV	ml/m <sup>2</sup>	Surface Area	ml/kg	Body Weight	ml/m <sup>2</sup>	Surface Area	ml/kg	Body Weight	RCV	PV	TBV
Sawka et al 7	18-35	51Cr 125I Alb	1000	1685	2685	26	43	69	-	-	-	-	-	-
Wennesland et al 4	19-52	51Cr -	1089	-	-	27	-	-	-	-	-	-	-	-
Huff & Feller 3	16-66	51Cr -	-	-	-	28	-	-	-	-	24	-	-	-
Retzlaff et al 5	20-66	51Cr T-1824	1095	1631	2726	27	40	67	834	1399	2233	23	39	62
Piomelli et al 10	18-29	51Cr	-	-	-	30	-	68	-	-	-	-	-	-
Schmidt et al 8	70-94	- T-1824	-	1610	-	-	-	-	-	1480	-	-	-	-
Sklaroff 9	60-99	- 131I Alb	-	-	-	-	37	-	-	-	-	-	32	-
Piomelli et al 10	75-92	51Cr	-	-	-	26	-	57	-	-	-	-	-	-
Present Study	64-100	51Cr 125I Alb	889	1634	2626	23	42	65	725	1424	2144	19	38	58

## REFERENCES

1. Gibson JG, Peacock WC, Seligman AM, Sack T: Circulating red cell volume measured simultaneously by the radioactive iron and dye methods. *J Clin Invest* 25:838-847, 1946.
2. Brady LW, Cooper DY, Colodzin M, McClenathan JE, King ER, Williams R: Blood volume studies in normal humans. *Surg Gynec Obstet* 97:25-32, 1953
3. Huff R, Feller DD: Relation of circulating red cell volume to body density and obesity. *J Clin Invest* 35:1-10, 1956.
4. Wennesland R, Brown E, Hopper J, Hodges JL, Guttentag OE, Scott KG, Tucker IN, Bradley B: Red cell, plasma and blood volumes in healthy men measured by radiochromium ( $\text{Cr}^{51}$ ) cell tagging and hematocrit: Influence of age, somatotype, and habits of physical activity on the variance after regression of volume to height and weight combined. *J Clin Invest* 38:1065-1077, 1959.
5. Retzlaff JA, Tauxe WN, Kiely JM, Stroebe CF: Erythrocyte volume, plasma volume, and lean body mass in adult men and women. *Blood* 33:649-661, 1969.
6. Hurley PJ: Red cell and plasma volume. *J Nucl Med* 16:46-52, 1975.
7. Sawka MN, Young AJ, Pandolf KB, Dennis RC, Valeri CR: Erythrocyte, plasma and blood volume of healthy young men. *Medicine and Science in Sports and Exercise* (in press).
8. Schmidt LA, Iob V, Flotte CT, Hodgson PE, McMath M: Blood volume changes in the aged. *Surgery* 40:938-944, 1956.
9. Sklaroff DM: Isotopic determination of blood volume in the normal aged. *Amer J Roentgenol* 75:1082, 1956.
10. Piomelli S, Nathan DG, Cummins JF, Gardner FH: The relationship of total red cell volume to total body water in octogenarian males. *Blood* 19:89-98, 1962.
11. Valeri CR, Cooper AG, Pivacek LE: Limitations of measuring blood volume with iodinated  $^{125}\text{I}$  serum albumin. *Arch Int Med* 132:534-538, 1973.
12. Valeri CR, Altschule MD: Hypovolemic anemia of trauma: the missing blood syndrome. Boca Raton: **CRC Press, Inc.**, 1981.
13. Cordts PR, Lamorte WW, DelGuercio C, Niehoff J, Pivacek LE, Fisher JB, Dennis RC, Siebens H, Giorgio A, Valeri CR, Menzoian JO: Red cell deficits after major elective surgery: poor predictive value of hematocrit and hemodynamic parameters. *Surg Gynec & Obstet* (in press).

14. Lamprecht W, Trautschold I: Determination with hexokinase and glucose-6-phosphate dehydrogenase. In: Bergmeyer HU, ed. Methods of enzymatic analysis. New York: Academic Press 1963:543-58.
15. Keitt AS: Pyruvate kinase deficiency and related disorders of red cell glycolysis. *Am J Med* 41:762-85, 1966.
16. Dennis RC, Bechthold D, Valeri CR: *In vitro* measurements of p50 - the pH correction and use of frozen red blood cells as controls. *Crit Care Med* 7:385-390, 1979.
17. Rodriguez JM, Shahidi NT: Erythrocyte 2,3-diphosphoglycerate in adaptive red cell volume deficiency. *NEJM* 285:479-482, 1971.